

Statistical analysis plan (SAP) for:

One-year clinical course of back-related disability and prognostic value of comorbidity on disability during one year follow-up in older patients visiting primary care with a new episode of back pain

Project:

BACK pain in Elders in Norway (BACE-N): a prospective cohort study of older people visiting primary care with a new episode of back pain

NCT identifier: NCT04261309
Document date: 22.03.2021

Table of Contents

| | |
|---|-------------------------------------|
| Authors:..... | Error! Bookmark not defined. |
| Project: | 1 |
| Administrative information | 3 |
| Scope | 3 |
| Working title..... | 3 |
| Version of SAP..... | 3 |
| Ethical approval..... | 3 |
| Names, affiliations and roles of SAP contributors | 3 |
| Study sponsor | 4 |
| Signatures | 4 |
| Introduction | 5 |
| Background and rationale for study | 5 |
| Study design and method | 5 |
| Study design | 5 |
| Study population..... | 5 |
| Data collection..... | 6 |
| Variables..... | 6 |
| Statistical power consideration..... | 7 |
| Statistical analyses..... | 8 |
| General statistical considerations | 8 |
| Statistical analyses..... | 8 |
| Handling of missing data..... | 7 |
| Sensitivity analyses | 9 |
| Proposed tables and figures | 10 |
| References..... | 15 |

Administrative information

Scope

This document is a supplement to the BACE-N protocol (ClinicalTrials.gov Identifier: NCT04261309). The current Statistical Analysis Plan has been written while data collection was ongoing (we had access to baseline data, but not to follow-up data) and it will be uploaded to the ClinicalTrials.gov before full access to the study database.

Working title

One-year clinical course of back-related disability and prognostic value of comorbidity on disability during one year follow-up in older patients visiting primary care with a new episode of back pain

Version of SAP

1.0

Ethical approval

The BACE-N study was deemed a “quality control project” by the Regional Ethical committee, and treatment by the ethical committee was thus not considered necessary as per 11.11.2014 (reference number: 2014/1634/REK vest)

Approval from the Norwegian Social Science Data Service was obtained on 02.03.2015 (reference number: 42149).

Names, affiliations and roles of SAP contributors

| Name | Title | Affiliation | Role |
|----------------------|--------------------|--|------------------------|
| Margreth Grotle | PT, professor, PhD | Oslo Metropolitan University P44, 0167 Oslo Phone: 90111172 Email: mgrotle@oslomet.no | Principal Investigator |
| Ørjan Nesse Vigdal | PT, PhD-student | Oslo Metropolitan University | Main author of SAP |
| Rikke Munk Killingmo | PT, PhD-student | Oslo Metropolitan University | Contributor to SAP |
| Kjersti Storheim | PT, professor, PhD | Oslo Metropolitan University | Contributor to SAP |
| Milada C. Småstuen | Professor, PhD | Oslo Metropolitan University | Statistical advisor |
| Lise Kretz | Chiro, PhD-student | Oslo Metropolitan University | Contributor to SAP |

Study sponsor

The BACE-N-study has received funding from Oslo Metropolitan University, The Norwegian Fund for Post-Graduate Training in Physiotherapy and “Et liv i bevegelse” (A life in movement) – Norwegian chiropractors’ research foundation.

Signatures

Ørjan Nesse Vigdal

Main author of SAP

Milada C. Småstuen

Statistical advisor

Margreth Grotle

Principal investigator

Introduction

Background and rationale for study

Back pain is common in all age groups (1), and one systematic review highlights that disabling back pain is more prevalent in older people than in younger people (2). The clinical course of back-related disability in older adults with back pain has not been extensively studied. Two studies suggests that improvements in disability were modest the first three months, with little to no improvements on group level after three months (3, 4). It is well documented that number of comorbidities are associated with the clinical course of back-related disability in older adults, but the prognostic value of comorbidity is still highly uncertain (5-11).

Study aim:

The primary aim of this study is to examine the clinical course of back-related disability measured at baseline, 3, 6 and 12 months after a new episode of back pain. The secondary aim is to assess the prognostic value of number and severity of comorbidity at baseline for changes in back-related disability over one year of follow-up.

Study design and method

Study design

BACE-N is based on the previously published BACE study protocol from the BACE international consortium (12). The BACE-N protocol has been published (ClinicalTrials.gov Identifier: NCT04261309).

The BACE-N study is a prospective observational cohort study with a 2-year follow-up time. This study will use data from baseline, 3, 6 and 12 months of follow-up. Design of the study was made within the framework PROgnosis RESearch Strategy (PROGRESS), which is a framework for ensuring and enhancing the quality of prognostic studies (13). The primary aim is relating overall prognosis, and the secondary aim is relating confirmatory prognostic factor research (13, 14).

Study population

Patients aged ≥ 55 years visiting a general practitioner (GP), physiotherapist or chiropractor for a new episode of back pain are invited to participate in the study. Back pain is defined as pain located in the region from the top of the scapula to the sacrum, with or without radiating leg pain. An episode of back pain is defined as “new” if the patient has not received health care for the same back complaint during the last 6 months. The exclusion criteria were: Difficulty completing the study questionnaires due to language or cognitive difficulties, mobility impairments impeding the clinical examination (wheelchair-bound patients), had received healthcare for the same back complaint during the last 6 months (notwithstanding care initiated within the previous 4 weeks from time of baseline assessment).

Data collection

At baseline, patients receive a clinical examination and questionnaire. Patients are then given a follow-up questionnaire either through email or mail at 3-months, 6-months and 12-months follow-up.

Description of treatment received during follow-up

The study participants continue their health care in agreement with their healthcare provider regardless of inclusion in the study. This means that patients may receive education and advice, exercise therapy, massage, manipulations, mobilizations, pharmacological therapy, or additional diagnostic testing and referrals, all of which constitutes “usual” primary care (15). Treatment given is at the discretion of the healthcare provider and the patient.

Variables

Outcome measure:

Back-related disability, measured using the Norwegian, validated version of the Roland-Morris Disability Questionnaire (RMDQ) (16, 17). This is a questionnaire with 24 statements regarding abilities to perform ADL tasks, with a dichotomous yes/no answer. The answers are summed to a total score ranging from 0-24, where 0 indicates no disability and 24 indicates “maximum” disability. RMDQ has been found to measure several dimensions of back-related disability (18). We plan to use RMDQ as a continuous measure, as recommended by the PROGRESS framework (14). This will ensure easier comparability to similar studies, and easier inclusion in future meta-analyses.

RMDQ is measured at baseline, 3 months, 6 months and 12 months.

Prognostic factor measurement:

The prognostic factor of interest is comorbidity measured at baseline with a modified version of the Self-administered Comorbidity Questionnaire (SCQ) (19). The original questionnaire measures 13 pre-specified comorbidities, and 2 non-specified. The item “back pain” has been removed for this study and replaced with an additional non-specified item. Thus, the count of comorbidity ranges from 0-15. The diseases listed are: Heart disease, high blood pressure, lung disease, diabetes, ulcer or stomach disease, kidney disease, liver disease, anemia or other blood disease, cancer, depression, osteoarthritis, rheumatoid arthritis and up to 3 non-specified comorbidities. The SCQ measures comorbidities on three levels: 1) Do you have the problem? 2) Do you receive treatment for it? 3) Does it limit your activities? All levels are answered on a dichotomous yes/no level, and you only answer level 2 and 3 if you have answered “yes” on level 1. An individual can receive a maximum of 3 points for each medical condition: 1 point for the presence of a comorbidity, 1 point if they receive treatment for the condition, and 1 point if the condition limits their functioning. The maximum score for the full SCQ is thus 45 points.

We plan to use SCQ part 1 comorbidity count (0-15 scale), and the full SCQ (0-45) in separate models. Previous studies have found linear relationships between comorbidity and back-related disability (7, 11), and thus we plan to treat these variables as linear. Linearity with outcome will be assessed, and deviations will be handled appropriately.

Covariates:

Covariates are presented in table 1. They are chosen based on being “established” prognostic factors in the literature, having been utilized in similar studies previously (to enhance comparability), and for being readily available and easy to measure in clinical practice.

| Table 1: Covariates, measurement level and/or instrument, rationale for inclusion | | |
|--|---|--|
| Factor | Measurement level | Rationale |
| Age | Continuous, minimum age ≥ 55 | Standard covariate in previous studies (7, 8, 10), and in some studies reported to be associated with disability levels in older adults (4, 6, 20) |
| Sex | Dichotomous | Standard covariate in previous studies (7, 8, 10), and in some studies reported to be associated with disability levels in older adults (6, 20) |
| BMI | Measured continuous, divided into categories: <20 , 20-25, 25-30, <35 | Covariate in previous study (7), and found to be associated with disability in older adults (4, 21) |
| Back pain duration | Measured in days, but categorized to an ordinal scale of 3 categories: 0-6 weeks, 6-12 weeks, >12 weeks, similar to other BACE studies (22) | Covariate in previous study (8), and found to be associated with disability in older adults (4, 6, 11, 20) |
| Baseline disability | Continuous. Roland-Morris disability questionnaire, 0-24 scale. | Standard covariate, given the outcome is disability. Only applicable if not using mixed models. |
| Pain severity last week | Continuous. Numeric rating scale 0-10. | Associated with disability in one study in older adults (4). |
| Expectation of recovery within the next three months | 5-point ordinal Likert scale. From “I am fully recovered” to “I am worse than ever”. | Consistently associated with disability in older adults (4, 6, 11) |

Statistical power consideration

The published protocol for the BACE-N study estimates that we need a total of 450 patients included in the study. Allowing a 15% dropout rate at 12 months, we will have approximately 380 participants available at 12 months.

We used the power estimation tool in Stata to determine level of statistical power. With a sample size of 450, and an estimated r^2 of 0.30 for the full model of comorbidity adjusted for covariates, we have over 90% power to detect an r^2 -change of 0.017 or higher when adding comorbidity to the model.

Handling of missing data

Missing data will be handled with multiple imputation, using 5 imputations and 10 iterations unless the missing exceeds 30% and missing at random cannot be assumed. We will use the fully conditioned specification method, and regression estimation. For variables where we are unable to use regression estimation due to computational difficulties, predictive mean matching will be used (23).

Missing values for RMDQ will be handled by replacing missing items with the mean of the answered items for the individual, if less than 30% of the items are missing.

A mixed model, which we intend to use for the primary and secondary aim, does not require variable level imputation for the outcome measure. However, missing values on item level on

the RMDQ might still be an issue. To solve this, we will replace missing items for RMDQ with the mean of the answered items for the individual, if less than 7 (30%) of the 24 items are missing.

Statistical analyses

General statistical considerations

Analyses described in this statistical analysis plan are considered a priori analyses. Possible post-hoc exploratory analyses will be explicitly identified in the article. All analyses will be carried out by the first author using Stata version 16, under supervision from the principal investigator and the advisory statistician. The statistical tests will be 2-sided, and p-value will be reported. If the p-value is less than 0.05, the test is deemed statistically significant. 95% confidence intervals will be reported on point estimates.

Statistical analyses

Description of study flow and study sample:

The flow of participants will be reported with a flow chart. Reasons for exclusion and loss to follow-up will be provided where known. Descriptive data of the study sample will be reported using mean and standard deviation for normally distributed continuous variables, median and interquartile range for variables with skewed distribution, and with frequency and proportions for categorical variables. Normal distribution will be examined visually using histogram and QQ-plot, and statistically with the Kolmogorov-Smirnov test. Baseline characteristics will be presented in a table. See proposed tables and figures below.

Descriptive statistics will be used to present the mean and standard deviation if RMDQ is normally distributed, or median and interquartile range if RMDQ is not normally distributed, for each time point: Baseline, 3 months, 6 months, 12 months. This will also be presented graphically, similarly to van der Gaag et al (3). This graphical presentation will also be performed stratified for number of comorbidities.

A person may be a responder at one time-point and a non-responder at another. Therefore, an analysis of responders versus non-responders will be performed for each time point, using bivariate analysis for baseline characteristics (chi square test, Individual Samples T-test, or Mann Whitney U-test). Results from these analyses will be presented in text, and the table available in supplementary material.

Model choice and model building strategy:

According to the STRATOS initiative task force, a complete prespecifying of all aspects in model building is unrealistic in observational studies (24). Thus, the following will provide a framework for analysis and model building in this study, not a detailed recipe. Decisions regarding final choice of models will be made by the first author, the advisory statistician and the principal investigator.

Primary aim, clinical course of back-related disability:

Mixed models for repeated measures will be used to account for statistical dependencies. RMDQ is the dependent variable, and time and first contact health provider will be entered as fixed factors. The exact handling of time as a continuous or categorical variable depends on the distribution. Previous studies have found that the clinical course of a back episode is not linear over time (3, 25, 26). It is therefore reasonable to believe that time will have to be treated as a categorical variable, or by introducing a quadratic term, depending on its distribution. Choice of covariance matrix is dependent on data structure. Interaction between first contact health provider and time will be analysed.

Secondary aim, prognostic value of number and severity of comorbidities on the course of back-related disability:

We will fit separate models for count of comorbidity (SCQ part 1, 0-15 scale) and count and severity of comorbidity (full SCQ, 0-45). The steps are outlined below:

1. The univariate association between SCQ and RMDQ over time will be assessed with a mixed model for repeated measures. An interaction term for SCQ*time will be tested, and kept if statistically significant. We will present the crude regression estimates from this analysis.
2. A mixed model with all the covariates from Table 1 and RMDQ over time will be fitted. From this model, r^2 will be presented as a measure of prognostic value.
3. SCQ will be added to the model from step 2. The adjusted regression estimate, and the r^2 -change from step 2 to step 3 will be presented.

Previous studies have found a linear relationship between number and severity of comorbidity and disability (7, 27). Linearity between SCQ and RMDQ will be assessed with a scatter-plot. In case of non-linearity between SCQ and RMDQ comorbidity count and disability score, the following alternatives will be discussed (24, 28):

- Categorization of SCQ
- Transformation (logarithmic or cubic) of SCQ
- Fractional polynomials for SCQ
- Spline functions for SCQ

Sensitivity analyses

Complete case analyses will be performed to assess possible bias introduced by the multiple imputation procedures.

Proposed tables and figures




Figure: Flowchart of study participants

Baseline characteristics:

Table: Baseline characteristics

| Variable | n | Values |
|---|---|--------|
| <i>Sociodemographic variables</i> | | |
| Age, median (IQR) | | |
| Female, n (%) | | |
| Married or living with partner, n (%) | | |
| Education level high, n (%) | | |
| <i>General health variables</i> | | |
| Health-related quality of life (SF-36), median (range) | | |
| Physical component score | | |
| Mental component score | | |
| Hazardous alcohol consumption (AUDIT-C), n (%) | | |
| Smoking status, n (%) | | |
| Current smoker | | |
| Smoked previously | | |
| Never | | |
| Falls efficacy (FES-I), median (range) | | |
| <i>Back pain history and characteristics of current episode</i> | | |
| First healthcare provider | | |
| General practitioner | | |
| Physical therapist | | |
| Chiropractor | | |
| History of back pain, n (%) | | |
| Using pain medication, n (%) | | |
| Sleep problems weekly due to back pain, n (%) | | |
| Back pain (NRS 0-10) (figure?), median (range) | | |
| Disability (RMDQ 0-24) (figure?), median (range) | | |
| Duration of current episode, n (%) | | |
| 0 – 6 weeks | | |
| 6 weeks – 3 months | | |
| > 3 months | | |
| <i>Psychological variables</i> | | |
| Kinesiophobia (FABQ-PA 0-28), median (range) | | |
| Depression (CES-D 0-60), median (range) | | |
| Pain catastrophizing (PCS 0-52), median (range) | | |
| Back beliefs and attitudes (BBQ 9-45), median (range) | | |
| Expectations of back pain next 3 months (figure?), n (%) | | |
| Fully recovered | | |
| Much better | | |
| No change or worse | | |
| Psychosocial risk profile (SBT) | | |
| Low risk | | |
| Medium risk | | |
| High risk | | |
| <i>Clinical variables</i> | | |
| Pain with active movements of the back, n (%) | | |
| Positive radiculopathy diagnostic rule, n (%) | | |
| Two or more red flags, n (%) | | |
| Physical performance (BPS), median (range) | | |
| Functional mobility (TUG), median (range) | | |

SF-36, Short Form health survey 36; AUDIT-C, alcohol use disorders identification test, score ≥ 3 for women and ≥ 4 for men ; FES-I, Falls Efficacy Scale – International; NRS, numeric rating scale; RMDQ, Roland-Morris Disability Questionnaire; FABQ-PA, Fear-Avoidance Beliefs Questionnaire – Physical Activity; CES-D, Center for Epidemiological Studies – Depression; PCS, Pain Catastrophizing Scale; BBQ, Back Beliefs Questionnaire; SBT, Start Back Tool; BPS, Back Performance Scale; TUG, Timed Up-and-Go.

*Comorbidities measured with Self-administered Comorbidity Questionnaire and a question on osteoporosis.

Appendix table: Baseline characteristics for responders vs non-responders for each time-point.

| Variable | Responder 3 mo (n) | Non-responder 3 mo | Responder 6 mo (n) | Non- responder 6 mo (n) | Responder 12 mo (n=) | Non- responder 12 mo (n=) |
|--|-----------------------|-----------------------|-----------------------|-------------------------------|-------------------------|---------------------------------|
| <i>Sociodemographic variables</i> | | | | | | |
| Age, median (IQR) | | | | | | |
| Female, n (%) | | | | | | |
| Married or living with partner, n (%) | | | | | | |
| Education level high, n (%) | | | | | | |

General health variables

Health-related quality of life
(SF-36), median (range)

Physical component score

Mental component score

Hazardous alcohol
consumption (AUDIT-C), n
(%)

Smoking status, n (%)

Current smoker

Smoked previously

Never

Falls efficacy (FES-I), median
(range)

Back pain history and characteristics of current episode

First healthcare provider

General practitioner

Physical therapist

Chiropractor

History of back pain, n (%)

Using pain medication, n (%)

Sleep problems weekly due to
back pain, n (%)

Back pain (NRS 0-10), median
(range)

Disability (RMDQ 0-24),
median (range)

Duration of current episode, n
(%)

0 – 6 weeks

6 weeks – 3 months

> 3 months

Psychological variables

Kinesiophobia (FABQ-PA 0-
28), median (range)

Depression (CES-D 0-60),
median (range)

Pain catastrophizing (PCS 0-
52), median (range)

Back beliefs and attitudes
(BBQ 9-45), median (range)

Expectations of back pain next
3 months (figure?), n (%)

Fully recovered

Much better

No change or worse

Psychosocial risk profile (SBT)

Low risk

Medium risk

High risk

Clinical variables

Pain with active movements of
the back, n (%)

Positive radiculopathy
diagnostic rule, n (%)

Two or more red flags, n (%)

Physical performance (BPS),
median (range)

Functional mobility (TUG),
median (range)

SF-36, Short Form health survey 36; AUDIT-C, alcohol use disorders identification test, score ≥ 3 for women and ≥ 4 for men ; FES-I, Falls Efficacy Scale – International; NRS, numeric rating scale; RMDQ, Roland-Morris Disability Questionnaire; FABQ-PA, Fear-Avoidance Beliefs Questionnaire – Physical Activity; CES-D, Center for Epidemiological Studies – Depression; PCS, Pain Catastrophizing Scale; BBQ, Back Beliefs Questionnaire; SBT, Start Back Tool; BPS, Back Performance Scale; TUG, Timed Up-and-Go.

*Comorbidities measured with Self-administered Comorbidity Questionnaire and a question on osteoporosis.


Prevalence of each comorbidity:

Table: Prevalence of each comorbidity at baseline

| Comorbidity | N | % |
|--|---|---|
| Heart disease | | |
| High blood pressure | | |
| Lung disease | | |
| Diabetes | | |
| Ulcer or stomach disease | | |
| Kidney disease | | |
| Liver disease | | |
| Anemia or other blood disease | | |
| Cancer | | |
| Depression | | |
| Osteoarthritis | | |
| Rheumatoid arthritis | | |
| Osteoporosis* | | |
| (Other non-prespecified comorbidities....) | | |

All comorbidities measured by Self-Administered Comorbidity Questionnaire

Example figure presenting clinical course, from van der Gaag et al (3):



Proportion of number of comorbidities among patients at baseline, example from Rundell et al (29):




Figure 1. Proportion for number of additional pain sites among those with persistent back pain.

RMDQ middle value presented at each time point, presented based on number of comorbidities (29):

Table: Unadjusted RMDQ score at each timepoint in total cohort and by number of comorbidities

| | Baseline | 3 months | 6 months | 12 months |
|-----------------------------|-----------------|-----------------|-----------------|------------------|
| Total cohort | Xx | Xx | Xx | xx |
| <i>No. of comorbidities</i> | | | | |
| 0 | Xx | Xx | Xx | xx |
| 1 | | | | |
| 2 | | | | |
| 3 | | | | |
| 4 | | | | |
| 5 | | | | |
| 6 | | | | |

RMDQ = Roland-Morris Disability Questionnaire

Association between comorbidities and disability during 1-year follow-up:

Table: Estimates from **mixed model(XX)** of effect of comorbidities on RMDQ score during 1 year of follow-up

| | β | 95% CI | R² |
|-----------------------------|---------------------------|---------------|----------------------|
| Comorbidities~ | | | |
| Covariates | | | |
| Comorbidities w/covariates* | | | |

RMDQ = Roland-Morris Disability Questionnaire

~unadjusted effect estimate

*adjusted for time, age, gender, education level, smoking status, hazardous alcohol consumption, back pain duration, probable radicular leg pain, (recruitment profession?)

^adjusted for age, gender, education level, smoking status, hazardous alcohol consumption, back pain duration, probable radicular leg pain, (recruitment profession?)

References

1. Hartvigsen J, Hancock MJ, Kongsted A, Louw Q, Ferreira ML, Genevay S, et al. What low back pain is and why we need to pay attention. *The Lancet*. 2018;391(10137):2356-67.
2. Dionne CE, Dunn KM, Croft PR. Does back pain prevalence really decrease with increasing age? A systematic review. *Age and ageing*. 2006;35(3):229-34.
3. van der Gaag WH, Enthoven WTM, Luijsterburg PAJ, van Rijckevorsel-Scheele J, Bierma-Zeinstra SMA, Bohnen AM, et al. Natural History of Back Pain in Older Adults over Five Years. *The Journal of the American Board of Family Medicine*. 2019;32(6):781-9.
4. Jarvik JG, Gold LS, Tan K, Friedly JL, Nedeljkovic SS, Comstock BA, et al. Long-term outcomes of a large, prospective observational cohort of older adults with back pain. *Spine J*. 2018;18(9):1540-51.
5. Scheele J, Enthoven WT, Bierma-Zeinstra SM, Peul WC, van Tulder MW, Bohnen AM, et al. Course and prognosis of older back pain patients in general practice: a prospective cohort study. *Pain*. 2013;154(6):951-7.
6. Deyo RA, Bryan M, Comstock BA, Turner JA, Heagerty P, Friedly J, et al. Trajectories of symptoms and function in older adults with low back disorders. *Spine (Phila Pa 1976)*. 2015;40(17):1352-62.
7. Leopoldino AAO, Megale RZ, Diz JBM, Moreira BDS, Lustosa LP, Pereira LSM, et al. Influence of the number and severity of comorbidities in the course of acute non-specific low back pain in older adults: longitudinal results from the Back Complaints in the Elders (BACE-Brazil). *Age and Ageing*. 2020;49(1):96-101.
8. Rundell SD, Resnik L, Heagerty PJ, Kumar A, Jarvik JG. Comparing Performance of Comorbidity Indices in Predicting Functional Status, Health-Related Quality of Life, and Total Health Care Use in Older Adults With Back Pain. *J Orthop Sports Phys Ther*. 2019:1-18.
9. Rundell SD, Goode AP, Suri P, Heagerty PJ, Comstock BA, Friedly JL, et al. Effect of comorbid knee and hip osteoarthritis on longitudinal clinical and health care use outcomes in older adults with new visits for back pain. *Archives of physical medicine and rehabilitation*. 2017;98(1):43-50.
10. Rundell SD, Resnik L, Heagerty PJ, Kumar A, Jarvik JG. Performance of the Functional Comorbidity Index (FCI) in Prognostic Models for Risk Adjustment in Patients With Back Pain. *PM&R*. 2020.
11. van der Gaag WH, Chiarotto A, Heymans MW, Enthoven WTM, van Rijckevorsel-Scheele J, Bierma-Zeinstra SMA, et al. Developing clinical prediction models for non-recovery in older patients seeking care for back pain: the BACE prospective cohort study. *PAIN*. 2020;Articles in Press.
12. Scheele J, Luijsterburg PA, Ferreira ML, Maher CG, Pereira L, Peul WC, et al. Back complaints in the elders (BACE); design of cohort studies in primary care: an international consortium. *BMC Musculoskelet Disord*. 2011;12:193.
13. Hemingway H, Croft P, Perel P, Hayden JA, Abrams K, Timmis A, et al. Prognosis research strategy (PROGRESS) 1: a framework for researching clinical outcomes. *BMJ*. 2013;346:e5595.
14. Riley RD, Hayden JA, Steyerberg EW, Moons KG, Abrams K, Kyzas PA, et al. Prognosis Research Strategy (PROGRESS) 2: prognostic factor research. *PLoS Med*. 2013;10(2):e1001380.
15. Kamper SJ, Logan G, Copsey B, Thompson J, Machado GC, Abdel-Shaheed C, et al. What is usual care for low back pain? A systematic review of health care provided to patients with low back pain in family practice and emergency departments. *PAIN*. 2020;161(4):694-702.
16. Roland M, Morris R. A Study of the Natural History of Back Pain: Part I Development of a Reliable and Sensitive Measure of Disability in Low-Back Pain. *spine*. 1983;8(2):141-4.
17. Grotle M, Brox J, Vollestad N. Cross-cultural adaptation of the Norwegian versions of the Roland-Morris Disability Questionnaire and the Oswestry Disability Index. *Journal of Rehabilitation Medicine*. 2003;35(5):241-7.

18. Grotle M, Brox JJ, Vøllestad NK. Functional Status and Disability Questionnaires: What Do They Assess?: A Systematic Review of Back-Specific Outcome Questionnaires. *Spine*. 2005;30(1):130-40.
19. Sangha O, Stucki G, Liang MH, Fossel AH, Katz JN. The self-administered comorbidity questionnaire: A new method to assess comorbidity for clinical and health services research. *Arthritis & Rheumatism*. 2003;49(2):156-63.
20. Jarvik JG, Comstock BA, Heagerty PJ, Turner JA, Sullivan SD, Xu S, et al. Back pain in seniors: the back pain outcomes using longitudinal data (BOLD) cohort baseline data. *BMC Musculoskeletal Disorders*. 2014;15(1):1-23.
21. Jesus-Moraleida FR, Ferreira PH, Ferreira ML, Silva JP, Maher CG, Enthoven WT, et al. Back Complaints in the Elders in Brazil and the Netherlands: a cross-sectional comparison. *Age and ageing*. 2017;46(3):476-81.
22. Scheele J, Enthoven WT, Bierma-Zeinstra SM, Peul WC, van Tulder MW, Bohnen AM, et al. Characteristics of older patients with back pain in general practice: BACE cohort study. *Eur J Pain*. 2013;18(2):279-87.
23. Van Buuren S. Flexible imputation of missing data: CRC press; 2018.
24. Sauerbrei W, Perperoglou A, Schmid M, Abrahamowicz M, Becher H, Binder H, et al. State-of-the-art in selection of variables and functional forms in multivariable analysis--outstanding issues. *arXiv preprint arXiv:190700786*. 2019.
25. Itz CJ, Geurts JW, Kleef M, Nelemans P. Clinical course of non-specific low back pain: A systematic review of prospective cohort studies set in primary care. *European Journal of Pain*. 2013;17(1):5-15.
26. Costa LdCM, Maher CG, Hancock MJ, McAuley JH, Herbert RD, Costa LO. The prognosis of acute and persistent low-back pain: a meta-analysis. *Canadian Medical Association Journal*. 2012;184(11):E613-E24.
27. Bruusgaard D, Tschudi-Madsen H, Ihlebæk C, Kamaleri Y, Natvig B. Symptom load and functional status: results from the Ullensaker population study. *BMC Public Health*. 2012;12(1):1085.
28. Riley RD, Moons KG, Hayden JA, Sauerbrei W, Altman DG. Prognostic Factor Research. In: Riley RD, van der Windt DA, Croft P, Moons KG, editors. *Prognosis Research in Healthcare Concepts, Methods and Impact*. Oxford, United Kingdom: Oxford University Press; 2019.
29. Rundell SD, Patel KV, Krook MA, Heagerty PJ, Suri P, Friedly JL, et al. Multisite Pain Is Associated with Long-term Patient-Reported Outcomes in Older Adults with Persistent Back Pain. *Pain Med*. 2019.